


Talochlamys pusio (Linnaeus, 1758)

AphiaID: 236715

HUMPBACK SCALLOP

Animalia (Reino) > Mollusca (Filo) > Bivalvia (Classe) > Autobranchia (Subclasse) > Pteriomorpha (Infraclasse) > Pectinida (Ordem) > Pectinoidea (Superfamília) > Pectinidae (Família)



© Vasco Ferreira

Synonyms

Chlamys distorta (da Costa, 1778)
Crassadoma pusio (Linnaeus, 1758)
Hinnites distorta (da Costa, 1778)
Hinnites megadesma Coen, 1930
Ostrea miniata Born, 1778
Ostrea pusio Linnaeus, 1758
Ostrea sinuosa Gmelin, 1791
Pecten crotilus Reeve, 1853
Pecten distortus da Costa, 1778
Pecten irregularis Deshayes, 1832
Pecten isabellae MacGillivray, 1843

Pecten senticosus Jeffreys, 1879

Pecten spinosus T. Brown, 1827

Pecten spinosus T. Brown, 1827

References

basis of record Schiaparelli, S. (2006). Bivalvi, in: Revisione della Checklist della fauna marina italiana. , available online at <http://www.faunaitalia.it/checklist/> [details]

context source (Deepsea) Janssen R., Krylova E.M. (2014). Deep-sea fauna of European seas: An annotated species check-list of benthic invertebrates living deeper than 2000 m in the seas bordering Europe. Bivalvia. Invertebrate Zoology. Vol.11. No.1: 43-82 [in English]. [details]

Last update: 19 Jul. 2019