

Callithamniella tingitana (Schousboe ex Bornet) Feldmann-Mazoyer, 1938

AphiaID: 144525

Biota (Superdominio) > Plantae (Reino) > Biliphyta (Subreino) > Rhodophyta (Filo) > Eurhodophytina (Subdivisao) > Florideophyceae (Classe) > Rhodymeniophycidae (Subclasse) > Ceramiales (Ordem) > Ceramiaceae (Familia) > Ceramioideae (Subfamilia) > Dohnielleae (Tribo) > Callithamniella (Genero)

Synonyms

Callithamnion aegagrophila Funk, 1922

Callithamnion tingitana Schousboe ex Bornet, 1892

Grallatoria tingitana (Schousboe ex Bornet) I.A.Abbott, 1976

References

additional source Guiry, M.D. & Guiry, G.M. (2019). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. , available online at <http://www.algaebase.org> [details]

basis of record Guiry, M.D. (2001). Macroalgae of Rhodophycota, Phaeophycota, Chlorophycota, and two genera of Xanthophycota, in: Costello, M.J. et al. (Ed.) (2001). European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification. Collection Patrimoines Naturels, 50: pp. 20-38 [details]

additional source Fredericq, S., T. O. Cho, S. A. Earle, C. F. Gurgel, D. M. Kravesky, L. E. Mateo-Cid, A. C. Mendoza-González, J. N. Norris, and A. M. Suárez. 2009. Seaweeds of the Gulf of Mexico, Pp. 187-259 in Felder, D.L. and D.K. Camp (eds.), Gulf of Mexico-Origins, Waters, and Biota. I. Biodiversity, pp. 187-259. Texas A&M Univ. Press. [details]

context source (Bermuda) Schneider, C. W. (2003). An annotated checklist and bibliography of the marine macroalgae of the Bermuda Islands. Nova Hedwigia, 76(3-4): 275-361 [details]

Last update: 23 Jul. 2019