

Aglaothamnion tenuissimum (Bonnemaison) Feldmann-Mazoyer, 1941

AphiaID: 144501

Biota (Superdominio) > Plantae (Reino) > Biliphyta (Subreino) > Rhodophyta (Filo) > Eurhodophytina (Subdivisao) > Florideophyceae (Classe) > Rhodymeniophycidae (Subclasse) > Ceramiales (Ordem) > Callithamniaceae (Familia) > Callithamnioideae (Subfamilia) > Callithamnieae (Tribo) > Aglaothamnion (Genero)

Sinónimos

Aglaothamnion byssoides (Arnott ex Harvey) C.F.Boudouresque & M.M.Perret-Boudouresque, 1987

Aglaothamnion furcellariae (J.Agardh) Feldmann-Mazoyer, 1941

Callithamnion arachnoideum C.Agardh, 1828

Callithamnion arnottii Trevisan, 1845

Callithamnion byssoides Arnott ex Harvey, 1833

Callithamnion furcellariae J.Agardh, 1851

Callithamnion hiemale Kjellman, 1907

Callithamnion tenuissimum (Bonnemaison) Kützinger, 1849

Ceramium tenuissimum Bonnemaison, 1828

Seirospora byssoides (Arnott ex Harvey) De Toni, 1903

Seirospora byssoides (Arnott ex Harvey) De Toni, 1903

Referências

additional source Guiry, M.D. & Guiry, G.M. (2018). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. , available online at <http://www.algaebase.org> [details]

basis of record Guiry, M.D. (2001). Macroalgae of Rhodophycota, Phaeophycota, Chlorophycota, and two genera of Xanthophycota, in: Costello, M.J. et al. (Ed.) (2001). European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification. Collection Patrimoines Naturels, 50: pp. 20-38 [details]

additional source Dyntaxa. (2013). Swedish Taxonomic Database. Accessed at www.dyntaxa.se [15-01-2013]., available online at <http://www.dyntaxa.se> [details]

Última atualização: 30 Nov. 2017

