

# *Lithophyllum orbiculatum* (Foslie) Foslie, 1900

---

AphiaID: 145157

Biota (Superdomínio) > Plantae (Reino) > Biliphyta (Subreino) > Rhodophyta (Filo) > Eurhodophytina (Subdivisao) > Florideophyceae (Classe) > Corallinophycidae (Subclasse) > Corallinales (Ordem) > Corallineae (Subordem) > Lithophyllaceae (Familia) > Lithophylloideae (Subfamilia) > Lithophylleae (Tribo) > Lithophyllum (Genero)

## Sinónimos

---

*Crodelia orbiculata* (Foslie) Kylin, 1956  
*Goniolithon subtenellum* Foslie, 1899  
*Lithothamnion orbiculatum* Foslie, 1895  
*Lithothamnion subtenellum* (Foslie) Me.Lemoine, 1915  
*Pseudolithophyllum orbiculatum* (Foslie) Me.Lemoine, 1929

## Referências

---

additional source Guiry, M.D. & Guiry, G.M. (2017). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. , available online at <http://www.algaebase.org> [details]

basis of record Guiry, M.D. (2001). Macroalgae of Rhodophycota, Phaeophycota, Chlorophycota, and two genera of Xanthophycota, in: Costello, M.J. et al. (Ed.) (2001). European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification. Collection Patrimoines Naturels, 50: pp. 20-38 [details]

additional source Sears, J.R. (ed.). 1998. NEAS keys to the benthic marine algae of the northeastern coast of North America from Long Island Sound to the Strait of Belle Isle. Northeast Algal Society. 163 p. [details]

additional source South, G. R. and I. Tittley. 1986. A checklist and distributional index of the benthic marine algae of the North Atlantic Ocean. Huntsman Marine Laboratory. St. Andrews, New Brunswick. 76 p. [details]

Última atualização: 22 Ago. 2017