Tritia reticulata (Linnaeus, 1758)

AphiaID: 876821

NETTED DOG-WHELK

Biota (Superdominio) > Buccinoidea (Superfamilia) > Nassariinae (Subfamilia) > Tritia (Genero)



Vasco Ferreira Todos os direitos reservados



Vasco Ferreira Todos os direitos reservados



Vasco Ferreira - OMARE / Dez. 30 2018 Todos os direitos reservados



Vasco Ferreira - OMARE / Dez. 30 2018 Todos os direitos reservados



Vasco Ferreira - OMARE / Dez. 30 2018 Todos os direitos reservados



Vasco Ferreira Todos os direitos reservados



v_s_ - iNaturalist.org

Identity confusions:



Tritia incrassata
Búzio



Tritia varicosa
Búzio

Synonyms

Nassarius reticulatus (Linnaeus, 1758)

References

new combination reference Galindo, L. A.; Puillandre, N.; Utge, J.; Lozouet, P.; Bouchet, P. (2016). The phylogeny and systematics of the Nassariidae revisited (Gastropoda, Buccinoidea). Molecular Phylogenetics and Evolution. 99: 337-353., available online at https://doi.org/10.1016/j.ympev.2016.03.019 [details]

status source Galindo, L. A.; Puillandre, N.; Utge, J.; Lozouet, P.; Bouchet, P. (2016). The phylogeny and systematics of the Nassariidae revisited (Gastropoda, Buccinoidea). Molecular Phylogenetics and Evolution. 99: 337-353., available online at https://doi.org/10.1016/j.ympev.2016.03.019 [details]

context source (BeRMS 2020) Marine Biology Section, Ugent. Belgium. INRAM. Benthic fauna

monitoring- SSD – Belgian Science Policy., available online at http://www.vliz.be/projects/inram/imers.php. [details]

Last update: 12 Dec. 2017