

Actinothoe sphyrodetta (Gosse, 1858)


AphiaID: 100986

ANÉMONA

Biota (Superdomínio) > Animalia (Reino) > Cnidaria (Filo) > Anthozoa (Classe) > Hexacorallia (Subclasse) > Actiniaria (Ordem) > Enthemonae (Subordem) > Metridioidea (Superfamília) > Sagartiidae (Família) > Actinothoe (Gênero)



Vasco Ferreira Todos os direitos reservados



Vasco Ferreira Todos os direitos reservados



Vasco Ferreira Todos os direitos reservados



Vasco Ferreira 



Vasco Ferreira 



Vasco Ferreira / Jul. 08 2013 Todos os direitos reservados



Vasco Ferreira / Jul. 08 2013 Todos os direitos reservados



Vasco Ferreira / Jul. 08 2013 Todos os direitos reservados




Vasco Ferreira / Jun. 04 2013 Todos os direitos reservados

Características identificativas

- Coluna esbranquiçada ou acinzentada, com listras longitudinais.
- Tentáculos brancos, translúcidos.
- Disco branco, amarelo-vivo ou laranja com marcas pretas na base dos tentáculos.
- Até 120 tentáculos.

- Tentáculos afastados até 3 cm do disco.
- Base da coluna com até 2 cm de diâmetro.

Estatuto de Conservação



Sinónimos

Actinia candida Gosse, 1853
Actinothoe (Sagartia) sphyrodeta (Gosse, 1858)
Actinothoe sphyrodeta candida Gosse
Actinothoe sphyrodeta xanthopis Gosse
Sagartia (Actinothoe) sphyrodeta (Gosse, 1858)
Sagartia candida
Sagartia sphyrodeta Gosse, 1858
Sagartia sphyrodita
Thoe sphyrodeta (Gosse, 1858)

Referências

Ballerstedt, S. 2007. *Actinothoe sphyrodeta* Sandalled anemone. In Tyler-Walters H. and Hiscock K. (eds) *Marine Life Information Network: Biology and Sensitivity Key Information Reviews*, [on-line]. Plymouth: Marine Biological Association of the United Kingdom. Available from: <http://www.marlin.ac.uk/species/detail/1872>

additional source Hayward, P.J.; Ryland, J.S. (Ed.). (1990). *The marine fauna of the British Isles and North-West Europe: 1. Introduction and protozoans to arthropods*. Clarendon Press: Oxford, UK. ISBN 0-19-857356-1. 627 pp. [\[details\]](#)

additional source Fautin, Daphne G. (2013). *Hexacorallians of the World.*, available online at <http://hercules.kgs.ku.edu/Hexacoral/Anemone2/> [\[details\]](#)

basis of record van der Land, J.; den Hartog, J.H. (2001). *Actiniaria*, in: Costello, M.J. et al. (Ed.) (2001). *European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification*. Collection Patrimoines Naturels, 50: pp. 106-109 [\[details\]](#)

additional source Muller, Y. (2004). Faune et flore du littoral du Nord, du Pas-de-Calais et de la Belgique: inventaire. [Coastal fauna and flora of the Nord, Pas-de-Calais and Belgium: inventory]. Commission Régionale de Biologie Région Nord Pas-de-Calais: France. 307 pp., available online at <http://www.vliz.be/imisdocs/publications/145561.pdf> [details]

additional source Fautin, Daphne G. (2013). Hexacorallians of the World. , available online at <https://web.archive.org/web/20191018194704/http://hercules.kgs.ku.edu/Hexacoral/Anemone2/> [details]

additional source Robins, M.W., 1969. The marine flora and fauna of the Isles of Scilly. Cnidaria e Ctenophora. J. nat. Hist. 3 : 329-343. [details]

additional source Teissier, G., 1950. Inventaire de la faune marine de Roscoff. Cnidaies et cténaies. Trav. biol. Roscoff. n. ser. suppl. 1: 1-43. [details]

additional source Ates, R. M. L.; Dekker, R.; Faasse, M. A.; den Hartog, J. C. (1998). The occurrence of *Sagartia elegans* (Dalyell, 1848) (Anthozoa: Actiniaria) in the Netherlands. Zoologische Verhandelingen. 323: 263-276. [details]

additional source Ocaña, Oscar; Den Hartog, J.C. (2002). A catalogue of Actiniaria and Corallimorpharia from the Canary Islands and from Madeira, Arquipelago. Life and Marine Sciences. 19A: 33-54. [details]

additional source den Hartog J.C., Ates R.M.L. (2011). Actiniaria from Ria de Arosa, Galicia, northwestern Spain, in the Netherlands Centre for Biodiversity Naturalis, Leiden. Zoologische Mededelingen, Leiden. 85 (2): 11-53. [details]

additional source Manuel, R. L. (1981). British Anthozoa. London: Academic Press. 241 pp. (Synopses of the British Fauna). [details]

additional source Cutress, B. D. (1961). *Habrosanthus bathamae*, n. gen., n. sp. (Actiniaria: Sagartiidae) from New Zealand. In: Transactions of the Royal Society of New Zealand. 1: 95-101. [details]

additional source Carlgren, O. (1949). A survey of the Ptychodactiaria, Corallimorpharia and Actiniaria. Kungliga Svenska Vet- enskapsakademiens Handlingar. 1: 1-121. [details]

additional source Harms, J. (1993). Check list of species (algae, invertebrates and vertebrates) found in the vicinity of the island of Helgoland (North Sea, German Bight) – a review of recent records. Helgolaender Meeresuntersuchungen, 47(1): 1-34, available online at <http://link.springer.com/article/10.1007/BF02366182> [details]

additional source den Hartog, J. C. (1977). Notes on the little known sea anemone *Cataphellia brodricii* and on the closely allied *Hormathia coronata* and *Paraphellia expansa* (Actiniaria, Hormathiidae). Netherlands Journal of Zoology, 27(3): 237-244 [details]

additional source Manuel, R. L. (1977). A redescription of *Edwardsia beautempsii* and *E. timida* (Actiniaria: Edwardsiidae). Cahiers de Biologie Marine, 18, 483-497 [details]

additional source Nafilyan, Z. (1912). Matériaux pour la faune des Actinies des côtes de France. Les Actinies de Roscoff. Memoirs of the Society of Zoology of France, 25, 5-44 [\[details\]](#)

additional source den Hartog, J. C. (1970). Verslag betreffende onderzoek aan zeeanemonen te Plymouth gedurende het tijdvak tussen 3 juni 1969 en 27 augustus 1969. Verslag van de Gewone Vergadering van de Afdeling Natuurkunde Koninklijke Nederlandse Akademie van Wetenschappen, 79(6): 96-100 [\[details\]](#)

additional source Ates, R. M. L. (1982). De Viltkokeranemoon – *Cerianthus lloydii*. Het Aquarium Maandblad voor Aquarium-, Terrarium- en Insektariumkunde, 52(3): 80-84 [\[details\]](#)

additional source Williams, R. B. (1997). *Actinotheroe sphyrodeta* (Cnidaria, Actiniaria): The first records from Portugal and the Mediterranean Sea. Journal of the Marine Biological Association of the United Kingdom, 77, 245-248 [\[details\]](#)

additional source Ates, R. M. L. (1997). Bloemdieren: De Zeeanemonen En Hun Verwanten Van De Nederlandse Kust. C. Moerman. Zeeanjer., pp. 31. [\[details\]](#)

additional source Teissier, G. (1965). Inventaire de la faune marine de Roscoff. Station Biologique de Roscoff. Roscoff., pp. 62. [\[details\]](#)

additional source Daly, M.; Gusmão, L.; Reft, A. J.; Rodríguez, E. (2010). Phylogenetic signal in mitochondrial and nuclear markers in sea anemones (Cnidaria, Actiniaria). Integrative and Comparative Biology, 50(3): 371-388 [\[details\]](#)

additional source Ramil Blanco, F.; Pulpeiro, E. F. (1990). Inventario de los antozoos de Galicia. A checklist of the Anthozoa from Galicia. Boletín de la Real Sociedad Española de Historia Natural, 86(1-4): 17-30 [\[details\]](#)

additional source Hayward, P.J. & J.S. Ryland (Eds.). (1990). The marine fauna of the British Isles and North-West Europe: 1. Introduction and protozoans to arthropods. Clarendon Press: Oxford, UK. 627 pp. [\[details\]](#)

Última atualização: 08 Out. 2018